

The Boeing Company
Rocketdyne Propulsion & Power
6633 Canoga Avenue
P.O. Box 7922
Canoga Park, CA 91309-7922

June 2, 2003
In reply refer to 2003RC02057



Mr. Michael Lopez
U. S. Department of Energy
National Nuclear Security Administration
(NNSA) Service Center
1301 Clay Street, Suite 700 N
Oakland, CA 94612-5208

Subject: NESHAPs Report for 2002

Dear Mr. Lopez:

Enclosed is the National Emission Standards for Hazardous Air Pollutants (NESHAPs) Report for 2002 for the Department of Energy's (DOE) facility at the Santa Susana Field Laboratory (SSFL). The U.S. Environment Protection Agency (EPA) regulates airborne releases of radioactivity from DOE facilities under 40 CFR 61, Subpart H. This document reports the radiochemical analysis results of the effluent samples from all applicable emission sources. It also includes the off-site dose assessment results, which are compared against the EPA standards for compliance demonstration.

During 2002, the only applicable emission source at the DOE facility at SSFL was the operating exhaust stack at the Radioactive Materials Handling Facility (RMHF). The RMHF Pond, usually considered an area source when it is dry, contained water all year long except for a few days, and, therefore, the sediment in the pond was not considered a potential source for airborne releases.

The EPA limit for a DOE site is 10 mrem/yr, as specified in 40 CFR 61, Subpart H. The regulation also specifies that radiation exposure dose to the Maximally Exposed Individual (MEI) be calculated using the EPA's CAP88PC computer model. Using the EPA's methodology, the Effective Dose Equivalent to the MEI from the RMHF exhaust during 2002 was calculated to be 1.5×10^{-6} mrem (1.5×10^{-8} mSv) per year. This result indicated that the release from the RMHF was about 6 million times less than the regulatory limit.

This report includes the Certification Statement to be signed by M. E. Lee (or designee) for The Boeing Company, Rocketdyne Propulsion & Power and by you for

M. Lopez
2003RC02057
6/2/2003
Page 2

the DOE Site Closure Office. The Certification Statements are required for the final report.

If you have any questions or comments on this report, please contact Ning Liu at (818) 586-6262.

Sincerely,



M. E. Lee, Program Manager
DOE Site Closure

ML:NL:je

Enclosure: Radionuclide Air Emissions Annual Report

cc: S. Black, DOE/NNSA Service Center

SHEA-097565



M. Lopez
2003RC02057
6/2/2003
Page 1

DOEAIR02

**U. S. Department of Energy
Radionuclide Air Emissions Annual Report
(under Subpart H of 40 CFR Part 61)
Calendar Year 2002**

Site Name: Santa Susana Field Laboratory
(Prepared on May 1, 2003)

Operations Office Information

Office: National Nuclear Security Administration (NNSA) Service Center
Address: 1301 Clay Street, Room 700N
Oakland, CA 94612-5208

Contact: Steve Black
Phone: (510) 637-1595

Site Information

Operator: The Boeing Company
Rocketdyne Propulsion & Power
Address: 6633 Canoga Avenue, MC T-038
P. O. Box 7922,
Canoga Park, CA 91309-7922

Contact: Ning Liu
Phone: (818) 586-6262

Section I. Facility Information

Site Description

The Santa Susana Field Laboratory (SSFL) is located at the boundary of Ventura and Los Angeles Counties in southern California, as shown in Figure 1. The site consists of four administrative areas and undeveloped land, with a total area of approximately 2,850 acres. A broad range of energy related research and development (R&D) projects, including nuclear technologies, were conducted in Area IV of the site. All the nuclear R&D operations in Area IV ceased in 1988, and the subsequent efforts have been directed toward decontamination and decommissioning (D&D) of the former nuclear facilities. Area IV has an area of about 290 acres, and Figure 2-1 shows the arrangement of the site.

The climate at SSFL is generally dry, with variable winds. The site is situated between Simi Valley and San Fernando Valley, and there is no significant agricultural land use within 30 km (19 miles) radius. While the land immediately surrounding Area IV is undeveloped, suburban residential areas are at greater distances.

Source Description

There are three radiological facilities or buildings remaining in Area IV of the SSFL, as shown in Figure 3. The Radioactive Materials Handling Facility (RMHF) is currently used for processing, packaging, and temporary storage of radioactive waste materials, which are eventually shipped off-site to DOE approved disposal facilities. As a result of the waste handling operations at the RMHF, radioactive effluents were released to the atmosphere through a stack in 2002. The effluents were filtered and monitored before released into the atmosphere to ensure compliance with the National Emission Standards for Hazardous Air Pollutants (NESHAP) requirements. No radioactive liquid effluents were released from the facility.

Building 4059 is the former Systems for Nuclear Auxiliary Power (SNAP) reactor ground test facility. The D&D of this facility started in the early 1990s. After the phase I remediation was completed in 1998, the ventilation system in this building was no longer needed, and the two ventilation stacks were demolished in 2000. In 2002, no radiological work requiring ventilation was performed in the building, and no effluents were released to the atmosphere from this building.

Building 4024 housed two experimental reactor systems during the 1960s. After the project was terminated, all equipment and fuel were removed from the facility. A portable HEPA filtered ventilation system was set up in the building to support the decontamination of the Hot Laboratory concrete blocks. The ventilation system has not been in operation since the decontamination work was completed in 1999. In 2002, no effluents were released to the atmosphere from this building.

M. Lopez
2003RC02057
6/2/2003
Page 3

The RMHF Pond (Sump 614) is a collection sump for rainfall runoff from the RMHF. This is a potential area source, because radioactivity in the sediment may become airborne when the pond is dry and the sediment is exposed to air. During 2002, the RMHF sump was covered with water except for a few days (sediment was still wet during those days). Therefore, no radionuclides from this potential source were resuspended by wind for airborne releases.

In 2002, the only applicable radiological emission source for the DOE facility at SSFL was the operating exhaust stack at the RMHF. Air samples from the ventilation stack were analyzed for specific radionuclides, and the results were used for the dose assessment in this report.




Figure 1. Location of Santa Susana Field Laboratory

Subdivisions			
Owner	Jurisdiction	Acres	Subtotals
Rocketdyne	Rocketdyne-Area IV	289.9	2,399.3
	Rocketdyne	784.8	
	Rocketdyne	1,324.6	
	(Undeveloped land)		
Government	NASA (former AFP 57)	409.5	451.2
	NASA (former AFP 64)	41.7	
Total Acres			2,850.5




Figure 2. Santa Susana Field Laboratory Site Arrangement




Figure 3. Potential Source Locations in Area IV at Santa Susana Field Laboratory

Section II. Air Emission Data

Point Source

<u>Point Source</u>	<u>Type Control</u>	<u>Efficiency</u>	<u>Location of MEI</u>
RMHF	Pre- and HEPA filters	99.97+%	2867 m NW

<u>Point Source Radionuclides</u>	<u>Annual Release Quantity</u>	
	(Ci)	(Bq)
H-3	2.0E-06	7.5E+04
Co-60	5.3E-08	2.0E+03
Sr-90	6.5E-08	2.4E+03
Y-90 (in equilibrium with Sr-90)	6.5E-08	2.4E+03
Th-230	3.4E-08	1.3E+03
U-234	9.6E-09	3.6E+02
U-235	6.6E-09	2.4E+02
Pu-238	2.1E-08	7.9E+02
Am-241	1.1E-08	3.9E+02

Area (Non-Point) Source

N/A

Section III. Dose Assessments

Description of Dose Model

Radiation doses to the Maximally Exposed Individual (MEI) as well as the population in the surrounding area resulting from the emissions of the DOE facility at SSFL during 2002 are calculated using the EPA's CAP88-PC computer code. Site-specific meteorological data, such as wind speed, direction frequency, and stability, were developed by the NRC and Argonne National Laboratory for SSFL and used for the atmospheric dispersion calculation in CAP88-PC. Other input data, such as release terms, stack height, and exhaust air velocity, were physically measured to represent the site-specific situation for dose calculations.

Demonstration of compliance with the NESHAPs standard is based on the calculation of the maximum radiation exposure dose to an offsite individual located at a residence, school, business, or office in the vicinity of the SSFL. For this purpose, the nearest such locations have been identified by review of maps, aerial photographs, and direct observations. They are the residential area of Simi Valley, the Brandeis-Bardin Institute, the Santa Monica Mountains Conservancy Sage Ranch office, the residential area in Black Canyon, and the residential area in Bell Canyon. The location with the highest estimated annual dose is considered the location of the Maximally Exposed Individual (MEI).

The RMHF stack is the only emission source in 2002. The Effective Dose Equivalent to the MEI resulting from the emission is compared against the NESHAPs standard for the demonstration of compliance. The dose was calculated using CAP88-PC with site-specific input data.

Compliance Assessment


Location of Maximally Exposed Individual (MEI): Residence in Simi Valley, 2867 m NW of RMHF.


The Effective Dose Equivalent to the MEI from the RMHF exhaust during 2002 was 1.5×10^{-6} mrem (1.5×10^{-8} mSv) per year. The EPA limit for a DOE site is 10 mrem/yr, as specified in 40 CFR 61, Subpart H. This result indicates that the release from the RMHF is about 6 million times less than the regulatory limit.

M. Lopez
2003RC02057
6/2/2003
Page 9

Certification

I certify under penalty of law that I have personally examined and am familiar with the information submitted herein and based on my inquiry of those individuals immediately responsible for obtaining the information, I believe that the submitted information is true, accurate and complete. I am aware that there are significant penalties for submitting false information including the possibility of fine and imprisonment. (See, 18 U.S.C. 1001).

 Date:
M. E. Lee
DOE Site Closure Program Manager
The Boeing Company
Rocketdyne Propulsion & Power

 Date: 6/2/03
M. Lopez
DOE Site Closure Office
NNSA Service Center
U. S. Department of Energy

Supplemental Information

The collective Effective Dose Equivalent resulting from the DOE operations at SSFL during 2002 is calculated to be 4.5×10^{-4} person-rem (4.5×10^{-6} person-Sv).

The population doses were calculated using CAP88-PC in the "POPULATION" mode. The site-specific population distribution is estimated from the demographic survey performed by Claritas Inc. Claritas Inc, a leading demographic survey company, developed the demographic data around SSFL in 2000 based on the census data and modified by direct observations of nearby residential areas around the SSFL site.

No operations regulated by Subparts Q and T were conducted in 2002, nor were there any emissions of Rn-220 from sources containing U-232 and Th-232. There were no non-disposal/non-storage sources of Rn-222 emission.

Potential releases from the RMHF are so low that, even assuming absence of the HEPA filters, estimated doses would be below the level that requires continuous monitoring as prescribed in 40CFR61.93(b). However, as a good practice, continuous monitoring is still being performed in accordance with ANSI N13.1 standard (1989 version). The stack effluent at RMHF is continuously sampled, counted for gross alpha and beta activities weekly, and combined annually for radiochemical analysis. In addition, tritium level in the evaporator feed water is analyzed annually to determine tritium release. This water is collected from the decontamination room at the RMHF, such as the wash down water for cleaning tools and equipment. The water is then stored in an 8000-gallon tank before evaporated through the HEPA filtered stack. Because of the very low level of tritium (lower than drinking water standards) and the large capacity of the storage tank, it is reasonable to assume that the tritium level is constant throughout the year.

There are five continuous ambient air samplers throughout the SSFL site. The purpose of this monitoring is to ensure that there is no airborne radioactivity resulted from the ongoing decontamination and decommissioning (D&D) activities at the site. Air sampling filters are collected and counted for gross alpha and beta activities on a weekly basis. The weekly samples are combined (separately by location) annually for radiochemical analysis. As usual, the ambient air samples collected in 2002 have radionuclide concentrations far below the Derived Concentration Guide (DCG) values and are generally indistinguishable from offsite background levels. Because the quantities are so close to the detection limits, the variability in the measurements is primarily due to analytical and background variations. The QA program described by Appendix B, Method 114 are implemented for this low-level surveillance effort.

Clean Air Act Assessment Package - 1988

S Y N O P S I S R E P O R T

Non-Radon Individual Assessment
Apr 15, 2003 05:05 pm

Facility: RMHF
Address: SSFL, Top of Woolsey Canyon Road, Simi Hills
City: Chatsworth
State: CA Zip: 91311-

Source Category: DOE facility
Source Type: Stack
Emission Year: 2002

Comments: CAP88PC calculation for 2002 ASER and NESHAPs
Maximum Exposed Individual, RMHF Stack

Effective Dose Equivalent
(mrem/year)

1.52E-06

At This Location: 2867 Meters Northwest

Dataset Name: RMHF02IND
Dataset Date: Apr 15, 2003 05:05 pm
Wind File: C:\CAP88PC2\WINDFILES\SSFLNRC.WND

MAXIMALLY EXPOSED INDIVIDUAL

Location Of The Individual: 2867 Meters Northwest
Lifetime Fatal Cancer Risk: 1.33E-11

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Dose Equivalent (mrem/y)
GONADS	2.04E-07
BREAST	7.83E-08
R MAR	1.44E-06
LUNGS	5.07E-06
THYROID	7.98E-08
ENDOST	1.71E-05
RMNDR	5.26E-07
EFFEC	1.52E-06

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source	TOTAL
			#1 Ci/y	
CO-60	Y	1.00	5.3E-08	5.3E-08
SR-90	D	1.00	6.5E-08	6.5E-08
Y-90	Y	1.00	6.5E-08	6.5E-08
H-3	*	0.00	2.0E-06	2.0E-06
U-234	Y	1.00	9.6E-09	9.6E-09
TH-230	Y	1.00	3.4E-08	3.4E-08
U-235	Y	1.00	6.6E-09	6.6E-09
PU-238	Y	1.00	2.1E-08	2.1E-08
AM-241	W	1.00	1.1E-08	1.1E-08

SITE INFORMATION

Temperature:	17 degrees C
Precipitation:	85 cm/y
Mixing Height:	366 m

SOURCE INFORMATION

Source Number: 1

Stack Height (m): 40.
Diameter (m): 1.

Plume Rise
Momentum (m/s): 15.
(Exit Velocity)

AGRICULTURAL DATA

	Vegetable	Milk	Meat
	<hr/>	<hr/>	<hr/>
Fraction Home Produced:	0.020	0.000	0.000
Fraction From Assessment Area:	0.000	0.000	0.000
Fraction Imported:	0.980	1.000	1.000

Food Arrays were not generated for this run.
Default Values used.

DISTANCES (M) USED FOR MAXIMUM INDIVIDUAL ASSESSMENT

2867

Clean Air Act Assessment Package - 1988

D O S E A N D R I S K E Q U I V A L E N T S U M M A R I E S

Non-Radon Individual Assessment

Apr 15, 2003 05:05 pm

Facility: RMHF
Address: SSFL, Top of Woolsey Canyon Road, Simi Hills
City: Chatsworth
State: CA Zip: 91311-

Source Category: DOE facility
Source Type: Stack
Emission Year: 2002

Comments: CAP88PC calculation for 2002 ASER and NESHAPs
Maximum Exposed Individual, RMHF Stack

Dataset Name: RMHF02IND
Dataset Date: Apr 15, 2003 05:05 pm
Wind File: C:\CAP88PC2\WNDFILES\SSFLNRC.WND

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)
GONADS	2.04E-07
BREAST	7.83E-08
R MAR	1.44E-06
LUNGS	5.07E-06
THYROID	7.98E-08
ENDOST	1.71E-05
RMNDR	5.26E-07
EFFEC	1.52E-06

PATHWAY EFFECTIVE DOSE EQUIVALENT SUMMARY

Pathway	Selected Individual (mrem/y)
INGESTION	5.51E-09
INHALATION	1.45E-06
AIR IMMERSION	5.41E-12
GROUND SURFACE	5.84E-08
INTERNAL	1.46E-06
EXTERNAL	5.84E-08
TOTAL	1.52E-06

NUCLIDE EFFECTIVE DOSE EQUIVALENT SUMMARY

Nuclide	Selected Individual (mrem/y)
CO-60	5.49E-08
SR-90	1.65E-09
Y-90	3.86E-11
H-3	1.86E-11
U-234	7.97E-08
TH-230	5.37E-07
U-235	5.40E-08
PU-238	4.28E-07
AM-241	3.62E-07
TOTAL	1.52E-06

CANCER RISK SUMMARY

Cancer	Selected Individual Total Lifetime Fatal Cancer Risk
LEUKEMIA	1.34E-12
BONE	7.58E-13
THYROID	3.13E-14
BREAST	2.61E-13
LUNG	8.54E-12
STOMACH	1.65E-13
BOWEL	8.43E-14
LIVER	1.76E-12
PANCREAS	1.11E-13
URINARY	7.04E-14
OTHER	1.36E-13
TOTAL	1.33E-11

PATHWAY RISK SUMMARY

Pathway	Selected Individual Total Lifetime Fatal Cancer Risk
INGESTION	3.60E-14
INHALATION	1.18E-11
AIR IMMERSION	1.31E-16
GROUND SURFACE	1.41E-12
INTERNAL	1.18E-11
EXTERNAL	1.41E-12
TOTAL	1.33E-11

NUCLIDE RISK SUMMARY

Nuclide	Selected Individual Total Lifetime Fatal Cancer Risk
CO-60	1.34E-12
SR-90	2.80E-14
Y-90	1.36E-15
H-3	5.09E-16
U-234	1.06E-12
TH-230	4.44E-12
U-235	7.53E-13
PU-238	3.77E-12
AM-241	1.86E-12
TOTAL	1.33E-11

INDIVIDUAL EFFECTIVE DOSE EQUIVALENT RATE (mrem/y)
(All Radionuclides and Pathways)

Distance (m)

Direction 2867

N	2.4E-07
NNW	8.9E-07
NW	1.5E-06
WNW	8.5E-07
W	1.7E-07
WSW	3.4E-07
SW	4.0E-07
SSW	3.4E-07
S	2.8E-07
SSE	6.9E-07
SE	1.1E-06
ESE	6.3E-07
E	1.5E-07
ENE	1.9E-07
NE	2.3E-07
NNE	2.3E-07

INDIVIDUAL LIFETIME RISK (deaths)
(All Radionuclides and Pathways)

Distance (m)

Direction 2867

N	2.1E-12
NNW	7.8E-12
NW	1.3E-11
WNW	7.5E-12
W	1.5E-12
WSW	3.0E-12
SW	3.5E-12
SSW	3.0E-12
S	2.4E-12
SSE	6.0E-12
SE	9.5E-12
ESE	5.5E-12
E	1.3E-12
ENE	1.7E-12
NE	2.0E-12
NNE	2.1E-12

Clean Air Act Assessment Package - 1988

SYNOPSIS REPORT

Non-Radon Population Assessment
Apr 15, 2003 05:12 pm

Facility: RMHF
Address: SSFL, Top of Woolsey Canyon Road, Simi Hills
City: Chatsworth
State: CA Zip: 91311-

Source Category: DOE facility
Source Type: Stack
Emission Year: 2002

Comments: CAP88PC calculation for 2002 ASER and NESHAPs
Population Dose, RMHF stack

Effective Dose Equivalent
(mrem/year)

1.74E-06

At This Location: 2400 Meters Northwest

Dataset Name: RMHF02POP
Dataset Date: Apr 15, 2003 05:12 pm
Wind File: C:\CAP88PC2\WNDFILES\SSFLNRC.WND
Population File: C:\CAP88PC2\POPPFILES\SSFL2000.POP

MAXIMALLY EXPOSED INDIVIDUAL

Location Of The Individual: 2400 Meters Northwest
 Lifetime Fatal Cancer Risk: 1.52E-11

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)	Collective Population (person-rem/y)
GONADS	2.36E-07	5.68E-05
BREAST	9.21E-08	1.98E-05
R MAR	1.65E-06	4.25E-04
LUNGS	5.81E-06	1.51E-03
THYROID	9.38E-08	2.01E-05
ENDOST	1.96E-05	5.09E-03
RMNDR	6.04E-07	1.54E-04
EFFEC	1.74E-06	4.49E-04

FREQUENCY DISTRIBUTION OF LIFETIME FATAL CANCER RISKS

Risk Range	# of People	# of People in This Risk Range or Higher	Deaths/Year in This Risk Range	Deaths/Year in This Risk Range or Higher
1.0E+00 TO 1.0E-01	0	0	0.00E+00	0.00E+00
1.0E-01 TO 1.0E-02	0	0	0.00E+00	0.00E+00
1.0E-02 TO 1.0E-03	0	0	0.00E+00	0.00E+00
1.0E-03 TO 1.0E-04	0	0	0.00E+00	0.00E+00
1.0E-04 TO 1.0E-05	0	0	0.00E+00	0.00E+00
1.0E-05 TO 1.0E-06	0	0	0.00E+00	0.00E+00
LESS THAN 1.0E-06	10222462	10222462	5.47E-08	5.47E-08

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source	TOTAL
			#1 Ci/y	Ci/y
U-234	Y	1.00	9.6E-09	9.6E-09
CO-60	Y	1.00	5.3E-08	5.3E-08
SR-90	D	1.00	6.5E-08	6.5E-08
Y-90	Y	1.00	6.5E-08	6.5E-08
AM-241	W	1.00	1.1E-08	1.1E-08
H-3	*	0.00	2.0E-06	2.0E-06
PU-238	Y	1.00	2.1E-08	2.1E-08
TH-230	Y	1.00	3.4E-08	3.4E-08
U-235	Y	1.00	6.6E-09	6.6E-09

SITE INFORMATION

Temperature:	17 degrees C
Precipitation:	85 cm/y
Mixing Height:	366 m

SOURCE INFORMATION

Source Number: 1

Stack Height (m): 40.
Diameter (m): 1.

Plume Rise
Momentum (m/s): 15.
(Exit Velocity)

AGRICULTURAL DATA

	Vegetable	Milk	Meat
	<hr/>	<hr/>	<hr/>
Fraction Home Produced:	0.020	0.000	0.000
Fraction From Assessment Area:	0.000	0.000	0.000
Fraction Imported:	0.980	1.000	1.000

Beef Cattle Density:	8.81E-02
Milk Cattle Density:	2.85E-02
Land Fraction Cultivated for Vegetable Crops:	1.18E-02

POPULATION DATA

	Distance (m)						
Direction	800	2400	4000	5600	7200	8800	11200
N	0	38	2055	7029	7689	5	14
NNW	0	0	3007	8597	6743	354	0
NW	0	562	6296	9532	10120	244	5
WNW	0	0	0	8175	7549	7392	1050
W	0	0	0	325	1129	193	12026
WSW	0	0	0	6	852	728	9539
SW	0	0	0	3554	3079	3041	11396
SSW	0	0	0	0	4166	5694	6453
S	0	305	3	0	1475	2306	3630
SSE	0	0	560	0	16	2991	4712
SE	0	188	832	1364	8154	9003	22249
ESE	0	201	0	2299	9422	13191	52287
E	0	0	0	1597	3483	5769	38595
ENE	0	0	0	280	387	2380	18881
NE	0	109	347	8915	2187	0	130
NNE	0	0	814	8261	2947	0	65

Direction	Distance (m)					
	14400	19200	25600	34400	48000	68000
N	0	202	1029	5024	434	873
NNW	0	1702	65	0	0	6131
NW	456	141	14242	748	255	346
WNW	13029	15438	804	30019	14607	35307
W	18841	13897	31911	46004	275199	540
WSW	20912	28943	12132	4209	18643	2
SW	13649	4018	925	108	0	0
SSW	1432	624	7206	0	0	0
S	681	5306	347	0	0	0
SSE	1007	4202	2979	0	13532	67961
SE	15379	5759	39912	493989	1110827	1461829
ESE	42581	90394	125998	465362	1623379	1719302
E	43909	133591	289752	263927	270781	283206
ENE	15185	45485	101785	6592	3517	74108
NE	63	15985	43962	69867	6882	208339
NNE	0	828	17371	11522	3341	2719

Clean Air Act Assessment Package - 1988

D O S E A N D R I S K E Q U I V A L E N T S U M M A R I E S

Non-Radon Population Assessment

Apr 15, 2003 05:12 pm

Facility: RMHF
Address: SSFL, Top of Woolsey Canyon Road, Simi Hills
City: Chatsworth
State: CA Zip: 91311-

Source Category: DOE facility
Source Type: Stack
Emission Year: 2002

Comments: CAP88PC calculation for 2002 ASER and NESHAPs
Population Dose, RMHF stack

Dataset Name: RMHF02POP
Dataset Date: Apr 15, 2003 05:12 pm
Wind File: C:\CAP88PC2\WNDFILES\SSFLNRC.WND
Population File: C:\CAP88PC2\POPFILES\SSFL2000.POP

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)	Collective Population (person-rem/y)
GONADS	2.36E-07	5.68E-05
BREAST	9.21E-08	1.98E-05
R MAR	1.65E-06	4.25E-04
LUNGS	5.81E-06	1.51E-03
THYROID	9.38E-08	2.01E-05
ENDOST	1.96E-05	5.09E-03
RMNDR	6.04E-07	1.54E-04
EFPEC	1.74E-06	4.49E-04

PATHWAY EFFECTIVE DOSE EQUIVALENT SUMMARY

Pathway	Selected Individual (mrem/y)	Collective Population (person-rem/y)
INGESTION	6.51E-09	1.33E-06
INHALATION	1.66E-06	4.33E-04
AIR IMMERSION	6.19E-12	1.61E-09
GROUND SURFACE	6.91E-08	1.41E-05
INTERNAL	1.67E-06	4.35E-04
EXTERNAL	6.91E-08	1.41E-05
TOTAL	1.74E-06	4.49E-04

NUCLIDE EFFECTIVE DOSE EQUIVALENT SUMMARY

Nuclides	Selected Individual (mrem/y)	Collective Population (person-rem/y)
U-234	9.12E-08	2.37E-05
CO-60	6.49E-08	1.33E-05
SR-90	1.91E-09	4.50E-07
Y-90	4.39E-11	1.07E-08
AM-241	4.15E-07	1.08E-04
H-3	2.12E-11	8.84E-09
PU-238	4.90E-07	1.27E-04
TH-230	6.14E-07	1.60E-04
U-235	6.19E-08	1.59E-05
TOTAL	1.74E-06	4.49E-04

CANCER RISK SUMMARY

Cancer	Selected Individual Total Lifetime Fatal Cancer Risk	Total Collective Population Fatal Cancer Risk (Deaths/y)
LEUKEMIA	1.54E-12	5.49E-09
BONE	8.67E-13	3.18E-09
THYROID	3.70E-14	1.08E-10
BREAST	3.08E-13	9.02E-10
LUNG	9.78E-12	3.57E-08
STOMACH	1.95E-13	5.74E-10
BOWEL	9.94E-14	2.92E-10
LIVER	2.02E-12	7.28E-09
PANCREAS	1.31E-13	3.85E-10
URINARY	8.31E-14	2.44E-10
OTHER	1.60E-13	4.71E-10
TOTAL	1.52E-11	5.47E-08

PATHWAY RISK SUMMARY

Pathway	Selected Individual Total Lifetime Fatal Cancer Risk	Total Collective Population Fatal Cancer Risk (Deaths/y)
INGESTION	4.25E-14	1.23E-10
INHALATION	1.35E-11	4.97E-08
AIR IMMERSION	1.49E-16	5.50E-13
GROUND SURFACE	1.66E-12	4.80E-09
INTERNAL	1.36E-11	4.99E-08
EXTERNAL	1.66E-12	4.80E-09
TOTAL	1.52E-11	5.47E-08

PATHWAY GENETIC RISK SUMMARY
(Collective Population)

Pathway	Genetic Risk (person-rem/y)
INGESTION	2.46E-08
INHALATION	1.61E-06
AIR IMMERSION	1.59E-09
GROUND SURFACE	1.39E-05
INTERNAL	1.64E-06
EXTERNAL	1.39E-05
TOTAL	1.55E-05

NUCLIDE RISK SUMMARY

Nuclide	Selected Individual Total Lifetime Fatal Cancer Risk	Total Collective Population Fatal Cancer Risk (Deaths/y)
U-234	1.21E-12	4.46E-09
CO-60	1.59E-12	4.61E-09
SR-90	3.25E-14	1.08E-10
Y-90	1.55E-15	5.31E-12
AM-241	2.13E-12	7.83E-09
H-3	5.80E-16	3.41E-12
PU-238	4.31E-12	1.59E-08
TH-230	5.07E-12	1.87E-08
U-235	8.65E-13	3.11E-09
TOTAL	1.52E-11	5.47E-08

INDIVIDUAL EFFECTIVE DOSE EQUIVALENT RATE (mrem/y)
(All Radionuclides and Pathways)

Distance (m)							
Direction	800	2400	4000	5600	7200	8800	11200
N	0.0E+00	2.8E-07	1.9E-07	1.5E-07	1.3E-07	1.1E-07	9.1E-08
NNW	0.0E+00	0.0E+00	7.1E-07	5.4E-07	4.6E-07	3.9E-07	0.0E+00
NW	0.0E+00	1.7E-06	1.2E-06	9.2E-07	7.8E-07	6.6E-07	5.2E-07
WNW	0.0E+00	0.0E+00	0.0E+00	5.3E-07	4.5E-07	3.8E-07	3.0E-07
W	0.0E+00	0.0E+00	0.0E+00	1.3E-07	1.1E-07	9.9E-08	8.0E-08
WSW	0.0E+00	0.0E+00	0.0E+00	2.0E-07	1.7E-07	1.4E-07	1.1E-07
SW	0.0E+00	0.0E+00	0.0E+00	2.0E-07	1.7E-07	1.5E-07	1.2E-07
SSW	0.0E+00	0.0E+00	0.0E+00	0.0E+00	1.6E-07	1.3E-07	1.1E-07
S	0.0E+00	3.2E-07	2.2E-07	0.0E+00	1.5E-07	1.3E-07	1.0E-07
SSE	0.0E+00	0.0E+00	5.3E-07	0.0E+00	3.3E-07	2.8E-07	2.2E-07
SE	0.0E+00	1.3E-06	8.2E-07	6.0E-07	5.0E-07	4.2E-07	3.3E-07
ESE	0.0E+00	7.4E-07	0.0E+00	3.6E-07	3.0E-07	2.5E-07	2.0E-07
E	0.0E+00	0.0E+00	0.0E+00	9.9E-08	8.8E-08	7.7E-08	6.2E-08
ENE	0.0E+00	0.0E+00	0.0E+00	1.2E-07	1.0E-07	9.0E-08	7.3E-08
NE	0.0E+00	2.7E-07	1.8E-07	1.4E-07	1.2E-07	0.0E+00	8.3E-08
NNE	0.0E+00	0.0E+00	1.9E-07	1.5E-07	1.3E-07	0.0E+00	8.7E-08

Distance (m)						
Direction	14400	19200	25600	34400	48000	68000
N	0.0E+00	4.9E-08	3.3E-08	2.2E-08	1.3E-08	6.4E-09
NNW	0.0E+00	1.6E-07	1.1E-07	0.0E+00	0.0E+00	2.3E-08
NW	3.9E-07	2.8E-07	1.9E-07	1.3E-07	7.8E-08	3.9E-08
WNW	2.3E-07	1.6E-07	1.1E-07	7.3E-08	4.4E-08	2.2E-08
W	6.1E-08	4.3E-08	2.8E-08	1.9E-08	1.1E-08	5.1E-09
WSW	8.4E-08	6.0E-08	4.0E-08	2.7E-08	1.7E-08	9.0E-09
SW	8.8E-08	6.4E-08	4.5E-08	3.2E-08	0.0E+00	0.0E+00
SSW	8.1E-08	5.8E-08	4.0E-08	0.0E+00	0.0E+00	0.0E+00
S	7.5E-08	5.3E-08	3.6E-08	0.0E+00	0.0E+00	0.0E+00
SSE	1.6E-07	1.2E-07	8.0E-08	0.0E+00	3.5E-08	1.9E-08
SE	2.5E-07	1.8E-07	1.2E-07	8.5E-08	5.4E-08	3.0E-08
ESE	1.5E-07	1.1E-07	7.4E-08	5.1E-08	3.2E-08	1.7E-08
E	4.7E-08	3.3E-08	2.2E-08	1.5E-08	8.7E-09	4.1E-09
ENE	5.5E-08	3.9E-08	2.6E-08	1.8E-08	1.0E-08	5.1E-09
NE	6.3E-08	4.4E-08	3.0E-08	2.0E-08	1.2E-08	6.1E-09
NNE	0.0E+00	4.7E-08	3.2E-08	2.1E-08	1.3E-08	6.3E-09

COLLECTIVE EFFECTIVE DOSE EQUIVALENT (person rem/y)
(All Radionuclides and Pathways)

	Distance (m)						
Direction	800	2400	4000	5600	7200	8800	11200
N	0.0E+00	1.1E-08	3.9E-07	1.1E-06	1.0E-06	5.7E-10	1.3E-09
NNW	0.0E+00	0.0E+00	2.1E-06	4.7E-06	3.1E-06	1.4E-07	0.0E+00
NW	0.0E+00	9.8E-07	7.6E-06	8.8E-06	7.9E-06	1.6E-07	2.6E-09
WNW	0.0E+00	0.0E+00	0.0E+00	4.4E-06	3.4E-06	2.8E-06	3.2E-07
W	0.0E+00	0.0E+00	0.0E+00	4.1E-08	1.3E-07	1.9E-08	9.6E-07
WSW	0.0E+00	0.0E+00	0.0E+00	1.2E-09	1.4E-07	1.0E-07	1.1E-06
SW	0.0E+00	0.0E+00	0.0E+00	7.3E-07	5.3E-07	4.4E-07	1.3E-06
SSW	0.0E+00	0.0E+00	0.0E+00	0.0E+00	6.6E-07	7.7E-07	6.9E-07
S	0.0E+00	9.7E-08	6.6E-10	0.0E+00	2.2E-07	2.9E-07	3.6E-07
SSE	0.0E+00	0.0E+00	3.0E-07	0.0E+00	5.2E-09	8.2E-07	1.0E-06
SE	0.0E+00	2.4E-07	6.8E-07	8.2E-07	4.1E-06	3.8E-06	7.3E-06
ESE	0.0E+00	1.5E-07	0.0E+00	8.3E-07	2.8E-06	3.4E-06	1.1E-05
E	0.0E+00	0.0E+00	0.0E+00	1.6E-07	3.1E-07	4.5E-07	2.4E-06
ENE	0.0E+00	0.0E+00	0.0E+00	3.3E-08	4.0E-08	2.2E-07	1.4E-06
NE	0.0E+00	2.9E-08	6.2E-08	1.2E-06	2.6E-07	0.0E+00	1.1E-08
NNE	0.0E+00	0.0E+00	1.5E-07	1.2E-06	3.7E-07	0.0E+00	5.7E-09

Direction	Distance (m)					
	14400	19200	25600	34400	48000	68000
N	0.0E+00	9.9E-09	3.4E-08	1.1E-07	5.7E-09	5.6E-09
NNW	0.0E+00	2.8E-07	7.2E-09	0.0E+00	0.0E+00	1.4E-07
NW	1.8E-07	3.9E-08	2.7E-06	9.5E-08	2.0E-08	1.4E-08
WNW	3.0E-06	2.5E-06	8.7E-08	2.2E-06	6.5E-07	7.8E-07
W	1.1E-06	5.9E-07	9.1E-07	8.8E-07	3.1E-06	2.8E-09
WSW	1.8E-06	1.7E-06	4.9E-07	1.2E-07	3.2E-07	1.8E-11
SW	1.2E-06	2.6E-07	4.1E-08	3.4E-09	0.0E+00	0.0E+00
SSW	1.2E-07	3.6E-08	2.9E-07	0.0E+00	0.0E+00	0.0E+00
S	5.1E-08	2.8E-07	1.2E-08	0.0E+00	0.0E+00	0.0E+00
SSE	1.7E-07	4.9E-07	2.4E-07	0.0E+00	4.7E-07	1.3E-06
SE	3.8E-06	1.0E-06	4.9E-06	4.2E-05	6.0E-05	4.3E-05
ESE	6.5E-06	9.8E-06	9.3E-06	2.4E-05	5.2E-05	2.9E-05
E	2.1E-06	4.4E-06	6.5E-06	3.9E-06	2.3E-06	1.2E-06
ENE	8.3E-07	1.8E-06	2.7E-06	1.2E-07	3.7E-08	3.8E-07
NE	4.0E-09	7.1E-07	1.3E-06	1.4E-06	8.4E-08	1.3E-06
NNE	0.0E+00	3.9E-08	5.5E-07	2.5E-07	4.3E-08	1.7E-08

INDIVIDUAL LIFETIME RISK (deaths)
(All Radionuclides and Pathways)

Distance (m)							
Direction	800	2400	4000	5600	7200	8800	11200
N	0.0E+00	2.5E-12	1.7E-12	1.3E-12	1.1E-12	9.9E-13	7.9E-13
NNW	0.0E+00	0.0E+00	6.2E-12	4.7E-12	4.0E-12	3.4E-12	0.0E+00
NW	0.0E+00	1.5E-11	1.1E-11	8.0E-12	6.7E-12	5.7E-12	4.5E-12
WNW	0.0E+00	0.0E+00	0.0E+00	4.6E-12	3.9E-12	3.3E-12	2.6E-12
W	0.0E+00	0.0E+00	0.0E+00	1.1E-12	9.8E-13	8.6E-13	6.9E-13
WSW	0.0E+00	0.0E+00	0.0E+00	1.7E-12	1.5E-12	1.2E-12	9.6E-13
SW	0.0E+00	0.0E+00	0.0E+00	1.8E-12	1.5E-12	1.3E-12	1.0E-12
SSW	0.0E+00	0.0E+00	0.0E+00	0.0E+00	1.4E-12	1.2E-12	9.2E-13
S	0.0E+00	2.8E-12	1.9E-12	0.0E+00	1.3E-12	1.1E-12	8.6E-13
SSE	0.0E+00	0.0E+00	4.6E-12	0.0E+00	2.8E-12	2.4E-12	1.9E-12
SE	0.0E+00	1.1E-11	7.1E-12	5.2E-12	4.3E-12	3.6E-12	2.8E-12
ESE	0.0E+00	6.4E-12	0.0E+00	3.1E-12	2.6E-12	2.2E-12	1.7E-12
E	0.0E+00	0.0E+00	0.0E+00	8.7E-13	7.7E-13	6.7E-13	5.4E-13
ENE	0.0E+00	0.0E+00	0.0E+00	1.0E-12	9.0E-13	7.8E-13	6.3E-13
NE	0.0E+00	2.4E-12	1.6E-12	1.2E-12	1.0E-12	0.0E+00	7.2E-13
NNE	0.0E+00	0.0E+00	1.6E-12	1.3E-12	1.1E-12	0.0E+00	7.6E-13

Distance (m)						
Direction	14400	19200	25600	34400	48000	68000
N	0.0E+00	4.2E-13	2.8E-13	1.9E-13	1.1E-13	5.5E-14
NNW	0.0E+00	1.4E-12	9.5E-13	0.0E+00	0.0E+00	2.0E-13
NW	3.4E-12	2.4E-12	1.6E-12	1.1E-12	6.7E-13	3.4E-13
WNW	2.0E-12	1.4E-12	9.3E-13	6.3E-13	3.8E-13	1.9E-13
W	5.2E-13	3.7E-13	2.5E-13	1.7E-13	9.6E-14	4.5E-14
WSW	7.3E-13	5.1E-13	3.5E-13	2.4E-13	1.5E-13	7.8E-14
SW	7.6E-13	5.5E-13	3.9E-13	2.7E-13	0.0E+00	0.0E+00
SSW	7.0E-13	5.0E-13	3.4E-13	0.0E+00	0.0E+00	0.0E+00
S	6.5E-13	4.6E-13	3.1E-13	0.0E+00	0.0E+00	0.0E+00
SSE	1.4E-12	1.0E-12	6.9E-13	0.0E+00	3.0E-13	1.6E-13
SE	2.1E-12	1.5E-12	1.1E-12	7.4E-13	4.7E-13	2.5E-13
ESE	1.3E-12	9.3E-13	6.4E-13	4.4E-13	2.7E-13	1.5E-13
E	4.1E-13	2.9E-13	1.9E-13	1.3E-13	7.5E-14	3.5E-14
ENE	4.8E-13	3.4E-13	2.3E-13	1.5E-13	9.1E-14	4.4E-14
NE	5.4E-13	3.8E-13	2.6E-13	1.8E-13	1.1E-13	5.3E-14
NNE	0.0E+00	4.0E-13	2.7E-13	1.8E-13	1.1E-13	5.4E-14

COLLECTIVE FATAL CANCER RATE (deaths/y)
(All Radionuclides and Pathways)

Direction	Distance (m)						
	800	2400	4000	5600	7200	8800	11200
N	0.0E+00	1.4E-12	4.9E-11	1.3E-10	1.2E-10	7.0E-14	1.6E-13
NNW	0.0E+00	0.0E+00	2.6E-10	5.7E-10	3.8E-10	1.7E-11	0.0E+00
NW	0.0E+00	1.2E-10	9.4E-10	1.1E-09	9.6E-10	2.0E-11	3.2E-13
WNW	0.0E+00	0.0E+00	0.0E+00	5.3E-10	4.1E-10	3.4E-10	3.9E-11
W	0.0E+00	0.0E+00	0.0E+00	5.1E-12	1.6E-11	2.3E-12	1.2E-10
WSW	0.0E+00	0.0E+00	0.0E+00	1.5E-13	1.8E-11	1.3E-11	1.3E-10
SW	0.0E+00	0.0E+00	0.0E+00	8.9E-11	6.5E-11	5.4E-11	1.6E-10
SSW	0.0E+00	0.0E+00	0.0E+00	0.0E+00	8.1E-11	9.4E-11	8.4E-11
S	0.0E+00	1.2E-11	8.2E-14	0.0E+00	2.7E-11	3.5E-11	4.4E-11
SSE	0.0E+00	0.0E+00	3.6E-11	0.0E+00	6.4E-13	1.0E-10	1.2E-10
SE	0.0E+00	3.0E-11	8.3E-11	1.0E-10	4.9E-10	4.6E-10	8.9E-10
ESE	0.0E+00	1.8E-11	0.0E+00	1.0E-10	3.5E-10	4.1E-10	1.3E-09
E	0.0E+00	0.0E+00	0.0E+00	2.0E-11	3.8E-11	5.5E-11	3.0E-10
ENE	0.0E+00	0.0E+00	0.0E+00	4.1E-12	4.9E-12	2.6E-11	1.7E-10
NE	0.0E+00	3.7E-12	7.7E-12	1.5E-10	3.2E-11	0.0E+00	1.3E-12
NNE	0.0E+00	0.0E+00	1.9E-11	1.5E-10	4.6E-11	0.0E+00	6.9E-13

Direction	Distance (m)					
	14400	19200	25600	34400	48000	68000
N	0.0E+00	1.2E-12	4.1E-12	1.4E-11	7.0E-13	6.8E-13
NNW	0.0E+00	3.4E-11	8.7E-13	0.0E+00	0.0E+00	1.7E-11
NW	2.2E-11	4.7E-12	3.2E-10	1.2E-11	2.4E-12	1.6E-12
WNW	3.6E-10	3.0E-10	1.1E-11	2.7E-10	7.9E-11	9.5E-11
W	1.4E-10	7.2E-11	1.1E-10	1.1E-10	3.7E-10	3.4E-13
WSW	2.1E-10	2.1E-10	5.9E-11	1.4E-11	3.9E-11	2.2E-15
SW	1.5E-10	3.1E-11	5.1E-12	4.2E-13	0.0E+00	0.0E+00
SSW	1.4E-11	4.4E-12	3.5E-11	0.0E+00	0.0E+00	0.0E+00
S	6.2E-12	3.4E-11	1.5E-12	0.0E+00	0.0E+00	0.0E+00
SSE	2.0E-11	6.0E-11	2.9E-11	0.0E+00	5.7E-11	1.5E-10
SE	4.7E-10	1.3E-10	6.0E-10	5.1E-09	7.3E-09	5.3E-09
ESE	7.9E-10	1.2E-09	1.1E-09	2.9E-09	6.3E-09	3.6E-09
E	2.5E-10	5.4E-10	7.9E-10	4.8E-10	2.9E-10	1.4E-10
ENE	1.0E-10	2.2E-10	3.3E-10	1.4E-11	4.5E-12	4.6E-11
NE	4.8E-13	8.7E-11	1.6E-10	1.7E-10	1.0E-11	1.6E-10
NNE	0.0E+00	4.7E-12	6.7E-11	3.0E-11	5.2E-12	2.1E-12